



AMPI-DRY 5000

Composition

Each bottle contains:

Ampicillin – Natrium	5 g
----------------------	-----

Characteristics

Ampi-Dry 5000 is indicated in the acute phase of infections where there is a need for bactericide antibiotic.

Indications

Infections of airways, infections of the gastrointestinal tract, urogenital infections, skin-infections, secondary bacterial infections in viral diseases, mastitis, (poly)-arthritis, septicaemia, omphalitis and meningitis caused by an unknown pathogen.

Ampi-Dry can also be used as prophylaxis during operations.

Dosage and administration

Administration: I.V., or I.M.

Horse/Cattle: 3 g per 100 kg bodyweight, 2 – 3 times a day, during 3 consecutive days.
Ampi Dry needs to be diluted in 26,5 ml water for injection to a final concentration of 5 g per 30ml.
The solution cannot be kept.

Adverse Reactions

Hypersensitive reactions may occur.

Contraindications

Do not administer to animals with a hypersensitivity for beta-lactam antibiotics.

Withholding period

Horse : 5 days

Cattle : 5 days

Milk (cattle): 2 days

Storage

Store at 15 – 25°C. No longer than 36 months.

Use solution immediately.

Keep out of reach and sight of children.

Presentation

Box 12 x 5 gram Ampi-Dry 5000 and 26.5 ml Water for Injection.

Registration (International)

NL 9887 / UDD