



EKYFLOGYL

Composition

Each 100 ml contains:

| | |
|----------------------------|--------|
| Prednisolone acetate | 200 mg |
| Lidocaine HCL | 1 g |
| Dimethyl sulphoxide (DMSO) | 88 ml |

Characteristics

DMSO has great penetrating properties and is an anti-inflammatory. It is also used as a vehicle to take other products through the skin so they can reach the inflamed area. Prednisolone is a corticosteroid with anti-inflammatory properties and Lidocaine is an analgesic.

Indications

Musculoskeletal inflammation with a traumatic or infectious aetiology. Arthritis, peri-arthritis, arthrosis, osteitis, synovitis, tendinitis, sprains and strains.

Dosage and administration

Apply with brush or gauze on area to be treated twice daily. Do not rub. Continue treatment until problem has resolved but with a maximum of 12 days.

Precautions

Do not resume training before all symptoms have cleared.
Wear gloves when applying product, avoid skin contact.

Contraindications

Do not use on animals with known liver or kidney problems.

Withholding period

Do not use on animals intended for food.

Warnings

External use only.
Do not use other products on area to be treated during treatment period.

Storage

Store at 15 – 25° C.
Keep out of reach and sight of children.

Presentation

125 ml bottle

Registration (International)

A.M.M. 683 580.0