



## ENGEMYCIN

### Composition

Each ml contains:

Oxytetracycline	100mg
-----------------	-------

### Characteristics

Broad-spectrum injectable antibiotic with a primarily bacteriostatic effect against numerous Gram-positive and Gram-negative bacteria, rickettsia, mycoplasma and chlamydia

### Indications

For the treatment of infections caused by bacteria sensitive to Oxytetracycline in horses, cattle, sheep and pigs. Examples include infections occurring in cases of pneumonia and other respiratory tract infections, (endo)metritis, mastitis, urinary tract infections, septicaemia and secondary infections associated with viral diseases.

### Dosage and administration

Engemycin can be administered at one of two dose rates. See leaflet for full details.

SA (Shorter action) For a shorter duration of activity, a dose rate of 3 – 10mg/kg bw (depending on species and age) I.M. or I.V., provides a duration of action of 24 hours following a single injection.

PA (Prolonged action): For a prolonged duration of activity, a dose rate of 10 – 20mg/kg bw (depending on species and age) I.M. only, provides a duration of action of 48 hours following a single injection.

### Withholding Period

- SA : Milk: 6 milkings – (72 hours)  
: Meat: Cattle, sheep and pigs – 10 days, horses – 63 days
- PA : Milk: 10 milkings – (120 hours)  
: Meat: Cattle, sheep and pigs – 14 days, PA not recommended for horses.  
Not for use in bobby calves.

### Storage

Store at 15 – 25°C.  
Keep away from children.

### Presentation

Bottle of 100 or 250 ml.

### Registration (International)

ACVM No. A3308