



## ADEQUAN I.M.

### Composition

Each ml contains:

Polysulphated glycosaminoglycan (PSGAG)	100 mg/ml
---	-----------

### Characteristics

Anti-Rheumatic. Solution for injection of horses.

### Indications

Lameness due to traumatic or degenerative aseptic joint disease.

### Dosage and administration

One 5ml ampoule I.M. every 4 days, total treatment consists of 7 injections.

### Adverse Reactions

A transitory local reaction at the injection site can occur in rare instances.

### Precautions

Wear gloves to avoid contact with the skin. Avoid self-injection.

### Contraindications

Do not use in horses that have bleeding tendencies or that have a prolonged clotting time, or within 24 hours post-operative.

Do not use in carrying or lactating mares.

Do not use in horses with liver- or kidney disorders or horses that are hypersensitive to PSGAG's

### Withholding period

The Netherlands: 0 days

### Warnings

Adequan I.M can enhance the actions of anti-coagulants

After overdosing, the clotting time can be prolonged for up to 8 hours.

Simultaneous use of NSAID's increases the risk of gastro-intestinal bleeding.

### Storage

Store dark at 15 – 30°C. Protect from freezing.

Keep out of reach and sight of children

### Presentation

7 vials of 5ml each.

### Registration (International)

REG NL 09616