



## EQ-ZONA

### Composition

Contains per sachet of 5g

Phenylbutazone	1 g
----------------	-----

### Characteristics

Granules/powder for oral administration with anti-inflammatory and analgesic properties.

### Indications

In equine:

Inflammatory processes in horses, local inflammation of the locomotion system as; Arthritis, Tendinitis, Myositis, Synovitis, Bursitis, Chronic Laminitis etc. General trauma, or post-operative treatment of the locomotion system, and pain management/treatment in general.

### Dosage and administration

Adult horses.

Can be administered with the daily feed-intake (topdressing) or mixed with the feed.

Day 1:	4.4 mg/kg/bw	2 sachets every 12 hours
Day 2 – 5 :	2.2 mg/kg/bw	1 sachet every 12 hours
Day 6 and hereafter:	2.2 mg/kg/bw	1 sachet every 24 hours

### Contraindications

Do not use on horses intended for consumption

### Warnings

Do not use in animals younger than 6 weeks of age. If such use cannot be avoided, animals may require a reduced dosage and careful clinical management. It is not recommended to use EQ-Zona during gestation or lactation.

### Storage

Store at 15 – 30°C.

Keep away from children.

### Presentation

Carton box with 100 sachets of 5 g

### Registration (International)

1173 ESP