



TENSOLVET GEL

Composition

Contains per 100g:

Heparin sodium	50.000	IU
Hydroxyethyl salicylate	5	g
Levomenthol	0,50	g
Chlorophyllin (E 141ii)		

Indications

For the treatment of local inflammatory swellings and bruising, including tendonitis, tenosynovitis, bursitis and other acute inflammatory conditions of the musculo-skeletal system in the horse. Tensolvét also promotes the early reabsorption of haematoma and oedematous swelling resulting from such conditions. Tensolvét has been shown to increase the blood flow through the underlying skin.

Dosage and administration

Prior to application ensure that the area to be treated is clean and dry. Apply Tensolvét to the intact skin using fingertip pressure only. For maximum effect animals should be treated up to 4 times a day.

10 – 20g of gel is usually sufficient for each application (depending on the area to be covered). The treated area is best left uncovered, but may be dressed if desired. Treatment can be continued for up to 6 days if necessary.

Adverse reactions

Animals *may*, infrequently, experience a mild skin reaction (which includes hair loss and blisters) following use of this product. If this occurs any remaining product should be thoroughly washed off, product use discontinued, and veterinary attention sought.

Precautions

Special precautions for use in animals

For external use on intact skin only.

Discontinue treatment at least 48 hours prior to racing

Do not massage after application

Discontinue if signs of discomfort or swelling occur

Special precautions for the person administering the veterinary medicinal product to animals

Impervious gloves should be worn when applying this product. Avoid contact with the eyes. In case of accidental contact with the eye, the affected eye(s) should be washed with copious quantities of water and medical advice sought as necessary. In case of accidental ingestion, seek medical attention immediately.

Contraindications

Do not apply to broken skin.

Do not use this product in cases where known salicylate hypersensitivity exists.

Warnings

For external use on intact skin only.

Discontinue treatment at least 48 hours prior to racing.

Withholding period

Meat and offal: 0 days

Do not use in mares whose milk is intended for human consumption.

Storage

Do not store above 25°C.

Protect from light.

Keep out of sight and reach of children

Presentation

Bottle à 300g.

Registration (international)

ATC Vet Code: QM02AC99